



Rejuvenation through art and science

Treatment Instructions



What to expect after the CoolSculpting procedure:

- » Typically, there is minimal discomfort and recovery time after the CoolSculpting procedure. Most patients are able to return to their daily routine immediately after the procedure.
- » Many patients experience redness in the treated area for up to a few hours after the applicator is removed. In rare cases, it may persist for up to two weeks.
- » Many patients have minimal discomfort following the procedure; however, you may experience one or more of the following sensations: swelling, itching, tingling, numbness, tenderness to the touch, pain in the treated area, cramping, aching, bruising and/or skin sensitivity. Consult your practitioner if these conditions persist beyond two weeks, or worsen over time.
- » Following the procedure, a gradual reduction in the thickness of the fat layer will take place. You may start to see changes as early as three weeks after the CoolSculpting procedure, and you will experience the most dramatic results after one to three months. Your body will continue to naturally process the injured fat cells from your body for approximately four months after your procedure.
- » *In rare cases*, patients have reported darker skin color, hardness, discrete nodules, freeze burn, enlargement of the treated area, hernia or worsening of existing hernia following the CoolSculpting procedure. Surgical intervention may be required to correct the enlargement or hernia.
- » Wearing a compression garment (i.e., Spanx) over the treated area/s may assist with your skin contouring.

Next steps:

- » Weight gain will prevent you from appreciating your full results. Maintain a healthy diet and exercise routine after your procedure to continue appreciating your full results.
- » Schedule a follow-up clinical assessment at approximately 4-6 weeks to review your clinical results and discuss the option of additional treatments to achieve your desired outcomes.

Please call us at 414-964-1111 with any questions, or if your symptoms appear to worsen or last longer than two weeks.

✓ Additional Instructions: _____